

TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371

760-248P

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/104132

INTERNATIONAL APPLICATION NO.

PCT/JP97/03946

INTERNATIONAL FILING DATE

October 30, 1997

PRIORITY DATE CLAIMED

October 30, 1996

TITLE OF INVENTION

c DNA FRAGMENT OF CAUSATIVE GENE OF SPINOCEREBELLAR ATAXIA TYPE2

APPLICANT(S) FOR DO/EO/US

TSUJI, Shoji; SANPEI, Kazuhiro

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39 (1).
4. ☐ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☐ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☒ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☒ A translation of the International Application into English (35 U.S.C. 371(c)(3)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(2)).
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A FIRST preliminary amendment.
☒ A SECOND or SUBSEQUENT preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:
 - 1). International Search Report
 - 2). Substitute Sequence Listing Paper & Disk Copy 14 Pages

U.S. APPLICATION NO (if known, see 37 CFR 1.5) <div style="text-align: center;">NEW</div>	INTERNATIONAL APPLICATION NO <div style="text-align: center;">PCT/JP97/03946</div>	ATTORNEY'S DOCKET NUMBER <div style="text-align: center;">760-248P</div>
---	---	---

17. <input checked="" type="checkbox"/> The following fees are submitted: BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(5)): Search Report has been prepared by the EPO or JPO \$930.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) \$720.00 No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)).... \$790.00 Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$1,070.00 International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4). \$98.00 <div style="text-align: right;">ENTER APPROPRIATE BASIC FEE AMOUNT =</div>	<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <th style="text-align: left;">CALCULATIONS</th> <th style="text-align: left;">PTO USE ONLY</th> </tr> <tr> <td colspan="2" style="height: 150px;"></td> </tr> <tr> <td style="text-align: right;">\$ 930.00</td> <td></td> </tr> <tr> <td style="text-align: right;">\$ 0.00</td> <td></td> </tr> </table>	CALCULATIONS	PTO USE ONLY			\$ 930.00		\$ 0.00	
CALCULATIONS	PTO USE ONLY								
\$ 930.00									
\$ 0.00									

Surcharge of \$130.00 for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).				\$ 0.00	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE		
Total Claims	13 - 20 =	0	X \$22.00	\$ 0.00	
Independent Claims	1 - 3 =	0	X \$82.00	\$ 0.00	
MULTIPLE DEPENDENT CLAIM(S) (if applicable) YES			+ \$270.00	\$ 270.00	
TOTAL OF ABOVE CALCULATIONS =				\$ 1,200.00	
Reduction of 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed (Note 37 CFR 1.9, 1.27, 1.28).				\$ 0.00	
SUBTOTAL =				\$ 1,200.00	
Processing fee of \$130.00 for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).				\$ 0.00	
TOTAL NATIONAL FEE =				\$ 1,200.00	
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +				\$ 40.00	
TOTAL FEES ENCLOSED =				\$ 1,240.00	
				Amount to be: refunded	\$
				charged	\$

a. ☒ A check in the amount of \$ **1,240.00** to cover the above fees is enclosed.

b. ☐ Please charge my Deposit Account. No. _____ in the amount of \$ _____ to cover the above fees.
 A duplicate copy of this sheet is enclosed.

c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any
 overpayment to Deposit Account No. 02-2448. A duplicate copy of this sheet is enclosed.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

Send all correspondence to:
Birch, Stewart, Kolasch & Birch, LLP
P.O. Box 747
Falls Church, VA 22040-0747
(703)205-8000

SIGNATURE
 MURPHY, GERALD M., JR.
 NAME
 28,977
 REGISTRATION NUMBER

/aam June 30, 1998

09/101132

760-248P
12 Rec'd PCT/PTO 30 JUN 1998

IN THE U. S. PATENT AND TRADE MARK OFFICE

APPLICANT: Shoji TSUJI et al.
INT'L. APPLN. NO.: PCT/JP97/03946
SERIAL NO.: New GROUP:
FILED: June 30, 1998 EXAMINER:
FOR: cDNA FRAGMENT OF CAUSATIVE GENE OF SPINOCEREBELLAR
ATAXIA TYPE2

PRELIMINARY AMENDMENT

Honorable Commissioner of Patents
BOX PATENT APPLICATION
Washington, D.C. 20231

June 30, 1998

Sir:

The following Preliminary Amendment and Remarks are respectfully submitted in connection with the above-identified application.

IN THE SPECIFICATION:

Before line 1, insert --This application claims the benefit under 35 U.S.C. §371 of prior PCT International Application No. PCT/JP97/03946 which has an International filing date of October 30, 1997 which designated the United States of America, the entire contents of which are hereby incorporated by reference.--

R E M A R K S


The specification has been amended to provide a cross-reference to the previously filed International Application.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account 02-2448 for any additional fees required under 37 C.F.R. §1.16 or under 37 C.F.R. §1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH, BIRCH, LLP

By


Gerald M. MURPHY

Reg. No. 28,977

P.O. Box 747

Falls Church, VA 22040-0747

GMM/aam
(703) 205-8000

IN THE U.S. PATENT AND TRADEMARK OFFICE

Applicant(s): Shoji TSUJI, Kazuhiro SANPEI
Application No.: NEW Group No.:
Filed: June 30, 1998 Examiner:
For: cDNA FRAGMENT OF CAUSATIVE GENE OF SPINOCEREBELLAR
ATAXIA TYPE 2

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

June 30, 1998

Dear Sir:

Preliminary to the examination of the above-referenced NEW U.S. patent application, the following amendments and remarks are respectfully submitted.

IN THE SPECIFICATION:

Page 3

Line 5, change "SEQ ID NO: 1" to --SEQ ID NO:2--

Page 4

Line 11, after "sequence " insert --(SEQ ID NO:1)--

Page 5

Line 6, change "ID NO: 1" to --ID NO:2--

Line 13, change "SEQ ID NO: 1" to --SEQ ID NO:2--

Page 12

Line 3, after “(CAG)₅₆” insert --(SEQ ID NO:6)--

Line 5, after “CC-3’ ” insert --(SEQ ID NO:3)--

Line 7, after first occurrence of “3’ ” insert --(SEQ ID NO:4)--

Line 7, after “TG-3’ ” insert --(SEQ ID NO:5)--

Page 15

Line 9, after “AGC-3’ ” insert --(SEQ ID NO:7)--

Line 10, after “3’ ” insert --(SEQ ID NO:8)--

Page 16

Line 11, after “C-3’ ” insert --(SEQ ID NO:9)--

Line 20, after “ACC-3’ ” insert --(SEQ ID NO:10)--

Page 17

Line 3, after “C-3’ ” insert --(SEQ ID NO:11)--

Line 4, after “CG-3’ ” insert --(SEQ ID NO:12)--

Page 18

Line 9, after “C-3’ ” insert --(SEQ ID NO:13)--

Please replace pages 21-33 with the Substitute Sequence Listing enclosed herewith.
Please renumber the remaining pages of the Specification, beginning with the Claims
consecutively from page 35 of the Substitute Sequence Listing.

IN THE CLAIMS:

Claim 1, line 2: after “sequence ” insert --as--.

Remarks

Enclosed herewith in full compliance with 37 C.F.R. 1.821-1.825 is a Substitute Sequence Listing to be inserted into the specification as indicated above. The Substitute Sequence Listing in no way introduces new matter into the specification.

Also submitted herewith in full compliance with 37 C.F.R. 1.821-1.825 is a disk copy of the Substitute Sequence Listing. The disk copy of the Substitute Sequence Listing, file 760-248.SUB, is identical to the paper copy, except that it lacks formatting.

If necessary, the Commissioner is hereby authorized in this, concurrent, and future replies, to charge payment or credit any overpayment to Deposit Account No. 02-2448 for any additional fees requires under 37 C.F.R. § 1.16 or under 37 C.F.R. § 1.17; particularly, extension of time fees.

Respectfully submitted,

BIRCH, STEWART, KOLASCH & BIRCH, LLP

By 

Gerald M. Murphy, Jr.
Reg. No. 28,977

P.O. Box 747
Falls Church, VA. 22040-0747
(703) 205-8000

GMM/jrr
760-248P

Enclosures: Paper and Disk copy of Substitute Sequence Listing

DESCRIPTION

cDNA Fragment of Causative Gene of Spinocerebellar Ataxia
Type 2

TECHNICAL FIELD

5 The present invention relates to cDNA fragments of
the causative gene of spinocerebellar ataxia type 2
(hereinafter also referred to as "SCA2"), proteins
encoded thereby, antibodies corresponding to the proteins,
and antisense nucleic acids of the above-mentioned cDNA
10 fragments.

BACKGROUND ART

SCA2 is an autosomal dominant, neurodegenerative
disorder that affects the cerebellum and other areas of
the central nervous system.

15 It has recently been discovered that the causative
genes of 6 neurodegenerative diseases including
dentatorubral-pallidoluysian atrophy (DRPLA) have more
CAG repeats than the normal genes. That is, the numbers
of CAG repeats in the causative genes of the
20 neurodegenerative diseases are 37 to 100, while those in
the normal genes are less than 35.

It has been suggested that the causative gene of
SCA2 has an increased number of CAG repeats (Trottier, Y.
et al. *Nature*, 378, 403-406 (1995)). However, since the
25 causative gene of SCA2 has not been identified, and since
its nucleotide sequence has not been determined, SCA2
cannot be diagnosed by a genetic assay.

DISCLOSURE OF THE INVENTION

An object of the present invention is to provide a sequence-determined cDNA fragment of the causative gene of SCA2. Another object of the present invention is to provide a protein produced by the causative gene of SCA2. Still another object of the present invention is to provide an antibody specific to the above-mentioned protein, which antibody is useful for therapy and diagnosis of SCA2. Still another object of the present invention is to provide an antisense of the causative gene of SCA2, which is useful for therapy of SCA2.

The present inventors intensively studied to discover a *Tsp* E1 fragment with a size of 2.5 kb in which the number of CAG triplet is increased only in SCA2 patients, and partial sequence thereof was determined. Human cDNA library was screened using as probes the oligonucleotides that respectively hybridize with the regions between which the CAG triplet repeats are interposed, and a cDNA fragment which hybridizes with both of these two probes was cloned. Using this cDNA fragment as a probe, human cDNA library was screened and a plurality of cDNA fragments which hybridize with the probe were cloned. Sequencing the cDNA fragments revealed that these cDNA fragments overlap with each other. To sequence the 5'-end and 3'-end regions, RACE (rapid amplification of cDNA ends) was performed. Further, to sequence the 5'-end region, RT-PCR was

performed, thereby succeeding in sequencing the full length of the cDNA of the causative gene of SCA2.

That is, the present invention provides a nucleic acid fragment comprising a nucleic acid region encoding an amino acid sequence shown in SEQ ID NO: 1 (provided that the number of repeat units of Gln from the 166th to 188th amino acid varies between 15 and 100). The present invention also provides a protein having an amino acid sequence encoded by the nucleic acid fragment according to the present invention. The present invention further provides an antibody which undergoes antigen-antibody reaction with the above-mentioned protein. The present invention still further provides an antisense nucleic acid having a size of not less than 15 bp, which hybridizes with a mRNA transcribed from the nucleic acid fragment according to the present invention so as to inhibit translation thereof.

By the present invention, a sequence-determined cDNA fragment of the causative gene of SCA2 was provided. The protein encoded by the nucleic acid fragment according to the present invention may be used for therapy of SCA2 and may be used as an immunogen for preparing an antibody useful for therapy and diagnosis of SCA2. Further, since the nucleotide sequence of the causative gene of SCA2 was determined by the present invention, antisense to this gene may now be designed. Still further, by the present invention, a recombinant vector comprising the nucleic

acid fragment according to the present invention which is incorporated in an expression vector that can express a desired gene in human body, which recombinant vector can express the nucleic acid fragment in human body, as well as a method for expressing the nucleic acid fragment according to the present invention comprising introducing the recombinant vector into human body. Thus, the present invention is thought to largely contribute to the therapy and diagnosis of SCA2.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows the nucleotide sequence of the cDNA fragment according to the present invention together with the amino acid sequence encoded thereby, which nucleotide sequence was determined in the Examples of the present invention.

Fig. 2 shows the continuation of Fig. 1.

Fig. 3 shows the continuation of Fig. 2.

Fig. 4 shows the continuation of Fig. 3.

Fig. 5 is a pedigree chart of the SCA2 patients who donated the genomic DNAs used in the Examples.

Fig. 6 shows the sizes, positions and restriction sites of the genomic DNA fragments Tsp1 and Tsp2, and SCA2 cDNA obtained in the Examples of the present invention, as well as the size and position of each of the obtained cDNA fragment.

Fig. 7 shows the distribution of the numbers of the CAG repeat units in the normal and SCA2 genes, which were

measured by using the (CAG)₅₅ probe.

BEST MODE FOR CARRYING OUT THE INVENTION

As described above, the nucleic acid fragment according to the present invention comprises the nucleic acid region encoding the amino acid sequence shown in SEQ ID NO: 1 in the SEQUENCE LISTING, provided that the number of repeating units of Gln from the 166th to the 188th amino acid varies between 15 and 100. The number of this repeat units is 15 to 25 in normal individuals and 35 to 100 in SCA2 patients. As is well-known, due to degeneration, there are a plurality of codons encoding one amino acid, and any nucleic acids which encode the amino acid sequence shown in SEQ ID NO: 1 are included within the scope of the present invention. The nucleotide sequence actually determined in the Examples below is shown in SEQ ID NO:1 and in Figs. 1-4. The way how the nucleotide sequence was determined and the fact that the cDNA having this nucleotide sequence is the cDNA of the causative gene of SCA2 are detailed in the Examples below.

The nucleic acid fragment according to the present invention may be cloned by the method detailed in the Examples below. Further, since the nucleotide sequence of the nucleic acid fragment according to the present invention was determined by the present invention, the nucleic acid fragment may be cloned by utilizing amplification by PCR using human cDNA library as a

template, or by hybridization using the PCR product as a probe. In cases where it is difficult to amplify the nucleic acid fragment by a single PCR, the nucleic acid fragment may be divided into a plurality of regions and the PCR products may be ligated by a conventional method so as to clone the nucleic acid fragment.

By incorporating the nucleic acid fragment according to the present invention into the multicloning site of a commercially available expression vector, and by transforming host cells with the obtained recombinant vector, the protein encoded by the nucleic acid fragment may be produced. Host-vector systems for expressing an arbitrary gene in a host cell is well-known in the art and a number of host-vector systems are commercially available. Those skilled in the art may easily express the nucleic acid fragment according to the present invention so as to produce the protein using such a commercially available host-vector system. Such a well-known method is described in, for example, D. M. Glover, DNA Cloning Volume III a practical approach, 1987, IRL Press.

For example, according to a conventional method, the nucleic acid fragment according to the present invention may be amplified by the long PCR (LA PCR) (Hiroyuki MUKAI, PROTEIN, NUCLEIC ACID, ENZYME, Vol.41, No.5, 585-594, 1996) using primers in which restriction sites are introduced, and the amplified product is digested with a

restriction enzyme. The digested amplified product is then inserted into the multicloning site of a commercially available plasmid vector pGEX (PHARMACIA) and ligated to obtain a recombinant vector, followed by transforming *E. coli* DH5 α (GIBCO BRL) by the conventional calcium chloride method. Transformants are then selected based on the drug resistance (ampicillin resistance) and the protein encoded by the causative gene of SCA2 may be recovered from the transformants. In cases where this vector is used, the desired protein is obtained as a fusion protein with GST (Glutathion S-Transferase), so that the protein may easily be detected using a commercially available anti-GST antibody (PHARMACIA).

Since the active protein produced by the gene of normal individuals (the number of CAG repeats is 15 to 25) is thought to have the normal function, SCA2 may be treated or alleviated by administering the normal protein to SCA2 patients. However, since SCA2 is autosomal dominant, it is necessary to simultaneously block the causative gene of SCA2 of the patient with the antisense described below.

By immunizing an animal with the above-described protein or an antigenic fragment thereof, an antibody specific to the protein or the fragment thereof may be recovered from the animal by a conventional method. The antibody may be a polyclonal antibody or a monoclonal antibody, and the monoclonal antibody may be prepared by

a well-known method.

It is theoretically difficult to construct an antibody using the polyglutamine chain encoded by the CAG repeat, which antibody has varying reactivities with the polyglutamine chain depending only on the lengths of the polyglutamine chain. However, by using the full length protein which was clarified by the present invention, an antibody which recognizes the stereoscopic difference between the patient's protein and the normal protein may be prepared. Further, by virtue of the present invention, since an antibody may be prepared by using a region of the SCA2 product, which is common to the normal individuals and the patients, a control is available in any assay system using an antibody specific to the protein according to the present invention. Using such an antibody, a simple assay such as plate method may be attained by a conventional method.

An affinity column may be prepared by immobilizing the antibody according to the present invention on agarose gel (e.g., Sephadex (trademark) of PHARMACIA), polystyrene beads or the like and by packing the resultant in a column. By passing a body fluid (blood, serum, spinal fluid or the like) through the affinity column, the protein produced by the causative gene of SCA2 of the patient may be obtained. By eluting the protein and by measuring the molecular weight thereof, diagnosis of SCA2 may be attained. This is because that

the protein of a SCA2 patient has a molecular weight larger than that of the protein of a normal individual because the number of CAG repeat units of the SCA2 causative gene in the patient is larger than that of the gene in the normal individual.

Possible gene therapies include the method by which production of the abnormal SCA2 gene product is stopped and the method by which normal SCA2 gene product is introduced. The former method includes blocking of translation of mRNA using an antisense. The latter method includes administration of the SCA2 gene product prepared *in vitro*, and introduction of the nucleic acid fragment into the cells. Since in patients suffering from SCA2 caused by extension of the CAG repeat, the abnormal protein is dominant, it is thought that the desired effect may be obtained by simultaneously performing both of the above-mentioned methods. In this case, the antisense is designed such that it hybridizes with a region not affecting the activity of the SCA2 product, and the region is removed from the normal SCA2 gene to be introduced, thereby assuring that the antisense does not inhibit the production of the normal protein. By designing the antisense and the normal SCA2 gene as mentioned above, the antisense and the SCA2 gene may be incorporated in the same vector and the vector may be introduced into cells.

The antisense nucleic acid according to the present

invention hybridizes with the mRNA transcribed from the nucleic acid fragment according to the present invention, and has a size of preferably not less than 15 bp and not more than the full length of the coding region of the nucleic acid fragment, more preferably not less than 50 bp and not more than the full length of the coding region of the nucleic acid fragment. Although the antisense nucleic acid preferably has a nucleotide sequence completely complementary to the entire mRNA or a part thereof, transcribed from the nucleic acid fragment according to the present invention, those having homologies to the degree that they hybridize with the mRNA *in vivo* are within the scope of the present invention. By administering the antisense according to the present invention to an SCA2 patient, the causative gene of SCA2 may be blocked. By administering the above-mentioned protein from a healthy individual to the patient under this condition, SCA2 may be treated or alleviated. The dose of the antisense nucleic acid may be appropriately selected depending on the conditions of the patient, and may usually be 0.001 mmol to 1000 mmol per day per 1 kg of bodyweight.

Known methods for introducing the gene into the body include methods in which Retrovirus or Adenovirus is used as a vector (Yasuhiro SETOYUCHI, Experimental Medicine, Vol.12, No.15 (extra edition) 1994, pp.114-121; Hiromi KANEGAE et al., Experimental Medicine, Vol.12, No.15

(extra edition) 1994, pp.34-40), methods using liposomes, and methods using fusion liposomes (membrane-fused liposome and the like). Among these, since Adenovirus expresses the introduced gene without proliferation of the target cells, the method utilizing Adenovirus is most appropriate as the method for introducing the nucleic acid into nervous system. More particularly, E3 and a part of Ela of the DNA of wild type Adenovirus type 5 are removed and the nucleic acid in double-stranded form is introduced into the virus together with an expression unit such as a promoter. The recombinant virus vector is proliferated in 293 cells expressing Ela and Elb genes, originated from human fetal kidney to prepare a virus liquid, and the virus liquid is inoculated. The virus genome incorporated into the nucleus of the cell exists outside the chromosomes without replication. Since the virus genome is not lost by cell division in nerve cells, expression of normal SCA2 gene may be maintained for 1 to 3 months.

The present invention will now be described more concretely by way of examples thereof. It should be noted that the present invention is not restricted to the following examples.

Example 1 Preparation of (CAG)₅₅ Probe

A genomic DNA segment of DRPLA gene containing a CAG repeat with 55 repeat units was amplified from the genomic DNA of a patient with DRPLA (Koide, R. et al.,

Nature Genet., 6, 9-13 (1994)) and was subcloned into a plasmid vector, pT7Blue T(*p*-2093). The *p*-2093 plasmid contains the (CAG)₅₅ and the flanking sequences. That is, the plasmid contains the sequence of 5'-CAC CAC CAG CAA CAG CAA (CAG)₅₅ CAT CAC GGA AAC TCT GGG CC-3'. Using a pair of oligonucleotides 5'-CAC CAC CAG CAA CAG CAA CA-3' and 5'-biotin-GGC CCA GAG TTT CCG TGA TG-3', PCR was performed in a total volume of 16 μ l containing 10 mM Tris-HCl, pH8.3, 50 mM KCl, 1.5 mM MgCl₂, 2M N,N,N-trimethylglycine, 0.1 mM TTP, 0.1 mM dCTP, 0.1 mM dGTP, 9.25 MBq of [α -³²P]dATP (222 TBq/mmol), 0.5 μ M each of the two primers, 0.3 ng of plasmid DNA (*p*-2093) and 2.0 U of Taq DNA polymerase (Takara Shuzo, Kyoto, Japan). After an initial 2-min. denaturation at 94°C, PCR was performed for 30 cycles consisting of 1-min. denaturation at 94°C, 1-min. annealing at 54°C and 3-min. extension at 72°C, followed by a final extension at 72°C for 10 min.

A single-stranded (CAG)₅₅ probe was isolated using streptavidin-coated magnetic beads (Dynabeads M-280, Streptavidin; Dynal AS, Oslo, Norway) on which 20 μ l of streptavidin is coated. That is, after washing of the PCR products immobilized on the magnetic beads with 40 μ l of a solution containing 5 mM Tris-HCl (pH 7.5), 0.5 mM EDTA and 1 M NaCl, the non-biotinylated strand containing the radio-label was separated from the biotinylated strand by incubation in 50 μ l of 0.1 M NaOH for 10 min. The resultant supernatant was directly added to the

hybridization solution described below.

Incidentally, using the single-stranded (CAG)₅₅ probe prepared as described above, Southern blot analysis was carried out on the androgen receptor genes containing 9, 22, 43 and 51 CAG repeat units, respectively. As a result, the (CAG)₅₅ probe strongly hybridized with the genes having 43 and 51 CAG repeats units, respectively, but scarcely hybridized with the gene having 22 CAG repeat units, and did not hybridize at all with the gene having 9 CAG repeat units (K. Sanpei et al., Biochemical and Biophysical Research Communications, Vol.212, No.2, 1995, pp.341-346). Thus, by using this probe, hybridization may be selectively attained only with DNAs containing a number of (e.g., 35 or more) CAG repeat units if the hybridization conditions are appropriately selected.

(2) Determination of Nucleotide Sequence of SCA2 Gene

Fig. 5 shows a pedigree chart of SCA2 patients. In this pedigree chart, males are represented by squares and females are represented by circles. SCA2 patients are represented by black squares or circles, and unaffected persons are represented by white squares or circles.

High-molecular-weight genomic DNA (15 µg) was digested with 100 U of *TspEI* (Toyobo, Osaka, Japan), electrophoresed through 0.8% agarose gels and transferred to nitrocellulose membranes. The membranes were hybridized with the (CAG)₅₅ probe described above.

Hybridization was performed in a solution containing 2.75 x SSPE (1 x SSPE=150 mM NaCl, 10 mM NaH₂PO₄, 1 mM EDTA), 50% formamide, 5 x Denhardt's solution, 100 ng/ml of sheared salmon sperm DNA and the (CAG)₅₅ probe (6 x 10⁶ cpm/ml) at 62°C for 18 hours. After the hybridization, the membranes were washed with 1 x SSC (150 mM NaCl, 15 mM sodium citrate) containing 0.5% SDS at 65°C for 0.5 hours. The membranes were autoradiographed for 16 hours to Kodak Bio Max MS film at -70°C using an MS intensifying screen.

As a result, 2.5 kbp *TspEI* fragment hybridized with the probe was detected only in all of the SCA2 patients.

Genomic DNA (270 µg) from an SCA2 patient (individual 7 in Fig. 5) was digested by *TspEI* and subjected to agarose gel electrophoresis. Genomic DNA fragments including the 2.5 kb *TspEI* fragment were cloned into an *EcoRI*-cleaved λZAPII vector. The genomic library was screened using the (CAG)₅₅ probe under the hybridization condition described above. A genomic clone, *Tsp-1*, containing an expanded CAG repeat was isolated.

After removal of the probe, the above-described genomic library was screened again using the *Tsp-1* as a probe, which was labeled by the random priming. Hybridization was carried out in a solution containing 5 x SSC, 1 x Denhardt's solution, 10% dextran sulfate, 20 mM sodium phosphate, 400 µg/ml human placental DNA and the *Tsp-1* probe at 42°C for 18 hours. After the

hybridization, the membranes were washed finally in 0.1 x SSC - 0.1% SDS at 52°C for 0.5 hours. The membranes were autoradiographed for 24 hours to Kodak Bio Max MS films at -70°C using an MS intensifying screen. As a result, a genomic clone, *Tsp-2*, originated from a normal allele was isolated.

The *Sma*I-*Apa*I fragment (630 bp) of *Tsp2* was sequenced and oligonucleotides F-1 (5'-CCC TCA CCA TGT CGC TGA AGC-3') and R-1 (5'-CGA CGC TAG AAG GCC GCT G-3') were designed such that the CAG repeat units are sandwiched between the oligonucleotides (see Fig. 1). Using oligonucleotides F-1 and R-1 as probes, human procephalic cortex cDNA library (STRATAGENE) was screened. Hybridization was performed in a solution containing 6 x SSC, 10 x Denhardt's solution, 0.5% SDS, 0.05% sodium pyrophosphate, 100 ng/ml of sheared salmon sperm DNA and end-labeled oligonucleotide probes at 55°C for 18 hours. After the hybridization, the membrane was finally washed with 6 x SSC containing 0.5% SDS and 0.05% sodium pyrophosphate at 55°C for 0.5 hours. A cDNA clone *Fc1* with a size of 4.0 kb which hybridized with the both probes was obtained. The nucleotide sequences of *Fc1*, *Tsp1* and *Tsp2* were determined and compared. As a result, the nucleotide sequences in the vicinities of the CAG repeat units were identical except for the number of the CAG repeat units. Restriction maps of *Tsp1* and *Tsp2*, as well as the sizes and positions of *Fc1* and other

fragments hereinbelow described, are shown in Fig. 6.

Using Fc1 or a fragment isolated by the screening later

described as a probe, human cDNA libraries (human

procephalic cortex, human fetal brain, human brain and

5 brain stem) were screened to isolate cDNA clones Fc2,

Fb14, B4, C6 and C19 (see Fig. 6). To identify the 5'-

end of Fc1, 5'-RACE (Frohman, M.A. et al, Proc. Natl.

Acad. Sci. USA 85, 8998-9002 (1988)) was performed using

5'-RACE-Ready cDNA (Clonetech, Palo Alto, CA, USA).

10 Primer R-1 was used for the first PCR, and Primer R-2

(5'-CTT GCG GAC ATT GGC AGC C-3', see Fig. 1) was used

for the second PCR. In both PCRs, F-1 (see Fig. 1) was

used as the forward primer. The 5'-RACE product (5R1)

having the size of 350 bp was subcloned into pT7Blue T

15 vector (pT7Blue T-vector (5R1)). The identification of

5R1 was confirmed by the overlapping with the nucleotide

sequences of Fc1, Tsp1 and Tsp2. To identify the 3'-end

of the cDNA, 3'-RACE was performed using 1 µg of

poly(A)⁺mRNA extracted from human brain as a template and

20 Primer F-13 (5'-TTC TCT CAG CCA AAG CCT TCT ACT ACC-3',

see Fig. 3) as a primer. The obtained 3'-RACE product

(3R1) with a size of 1300 bp was subcloned into pT7Blue T

vector (pT7Blue T-vector (3R1)).

To investigate the 5'-end region of the cDNA,

25 reverse transcription PCR (RT-PCR) was performed. That

is, total RNAs extracted from an autopsy from human brain

were digested by RNase-free DNase (PROMEGA) (Onodera, O.

et al., Am. J. Hum. Geent. 57, 1050-1060(1995)). As the primers for the PCR, F1006 (5'-TAT CCG CAG CTC CGC TCC C-3', see Fig. 1) and R1002 (5'-AGC CGG GCC GAA ACG CGC CG-3') were used. PCR was performed in a solution with a total volume of 20 μ M, which contained 5 pmol each of the each primer, 10 mM Tris HCl (pH8.3), 50 mM KCl, 1.5 mM $MgCl_2$, 1.7M N,N,N-trimethylglycine, 200 μ M each of dATP, dCTP and TTP, 100 μ M of dGTP, 100 μ M of 7-deaza dGTP and 2.5 U of Taq polymerase (TAKARA SHUZO). After carrying out the initial denaturation at 96°C for 2 minutes, a cycle of a denaturation step at 96°C for 1 minute, an annealing step at 65°C for 1 minute and an extension step at 72°C for 1 minute were repeated 30 times, and a final extension step at 72°C for 5 minutes was performed, thereby carrying out the PCR. As a result, a clone 5R1 which extends upstream of 5R1 by 246 bp was obtained (see Fig. 6).

In Fig. 6, the hollow regions in the Tsp1 and Tsp2 fragments indicate the regions which exist in SCA2 cDNAs. The hollow regions in the SCA2 cDNA shows coding regions. The CAG repeating regions are shown as solid boxes. Restriction sites TspE1 (T) , NotI (N), Sac II (S), Sau3AI (Sa) , Sma I (Sm), Eco52I (E52), Apa I (Ap), AccI (Ac) , BamHI (B) , XhoI (X), EcoRI (E) and Pst I (P) are shown. The size and position of each cDNA clone are shown below the consensus SCA2 cDNA.

In this example, nucleotide sequences of double-

stranded DNAs were determined by the dideoxynucleotide chain termination method (Sanger, F. et al. Proc. Natl. Acad. Sci. USA 74, 5463-5467(1977); Chen E.Y. et al, DNA 4, 165-170 (1985)) using a double-stranded plasmid DNA as a template. To determine the nucleotide sequences of the CAG repeating regions and their flanking regions, genomic fragments containing the CAG repeating regions were amplified by PCR using biotinylated F-1 and RS-1 (5'-CCT CGG TGT CGC GGC GAC TTC C-3'). PCR was performed in a solution with a total volume of 25 µl, which contained 0.25 µM each of the each primer, 10 mM Tris HCl (pH8.3), 50 mM KCl, 2.0 mM MgCl₂, 1.7M N,N,N-trimethylglycine, 200 µM each of dNTP, 200 ng of the genomic DNA and 1.25 U of Taq polymerase (TAKARA SHUZO). After carrying out initial denaturation at 95°C for 1 minute, a cycle of a denaturation step at 95°C for 2 minutes, an annealing step at 62°C for 1 minute and an extension step at 72°C for 1 minute was repeated 32 times, and a final extension step at 72°C for 5 minutes was performed, thereby carrying out the PCR. Biotinylated chains were recovered using streptavidin-coated magnetic beads and were directly sequenced.

Based on the nucleotide sequences of the above-mentioned cDNA clones, a consensus SCA2 cDNA sequence with a length of 4351 bp excluding the poly A tail was determined (SEQ ID NO:1, Figs. 1-4, see Fig. 6). In SEQ ID NO: 1, the region from 4352nt to 4367nt is the poly A

tail, and the number of "A" is not restricted to that shown in SEQ ID NO: 1. It was confirmed that the poly A tail exists at the same location in C19, B4 and 3R1 which were independent cDNA clones.

5 Example 2 Measurement of CAG Repeat Units in Sample

Numbers of CAG repeat units were determined by polyacrylamide gel electrophoresis analysis of PCR products obtained using the primer pair of F-1 and R-1. PCR was performed in a total volume of 10 μ l containing 10 mM Tris-HCl, pH 8.3, 50 mM KCl, 2.0 mM MgCl₂, 1.7 M *N,N,N*-trimethylglycine, 111KBq of [α -³²P]dCTP (111 Tbq/mmol), 30 μ M dCTP, and 200 μ M each of dATP, dGTP and TTP, 0.25 μ M each of the two primers, 200 ng of genomic DNA and 1.25 U of Taq DNA polymerase. After an initial 2-min denaturation at 95°C, PCR was performed for 32 cycles of 1-min denaturation at 95°C, 1-min annealing at 60°C and 1-min extension at 72°C, followed by a final extension at 72°C for 5 min. Sequence ladders obtained using the cloned genomic segments of the SCA2 gene, which contain various sizes of CAG repeats, were used as size markers. For normal alleles containing one or two CAA interruptions, the numbers of the CAA units were included in the CAG repeat size. For SCA2 alleles having expanded CAG region, the above-mentioned insert sequence immediately after the CAG region was not included in the size of the CAG region.

By the above-described method, the numbers of the

CAG repeat units of normal individuals (286 chromosomes) and 10 pedigrees of SCA2 patients (34 SCA2 chromosomes) were determined. The results are shown in Fig. 7. In Fig. 7, open bars indicate the results of the normal genes and solid bars indicate the results of the SCA2 genes.

As is apparent from Fig. 7, in all of the normal genes, the numbers of the CAG repeat units were not more than 24, while in all of the SCA2 genes, they were not less than 35. Thus, it was confirmed that the cDNA identified as described above is the cDNA of the causative gene of SCA2.

SEQUENCE LISTING

SEQ ID NO: 1

SEQUENCE LENGTH: 4367

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: double

TOPOLOGY: linear

SEQUENCE DESCRIPTION

TATCCGCACC TCCGCTCCCA CCCGGCGCCT CGGCGCGCCC GCCCTCCG ATG CGC TCA	57
Met Arg Ser	
1	
GCG GCC GCA GCT CCT CGG AGT CCC GCG GTG GCC ACC GAG TCT CGC CGC	105
Ala Ala Ala Ala Pro Arg Ser Pro Ala Val Ala Thr Glu Ser Arg Arg	
5 10 15	
TTC GCC GCA GCC AGG TGG CCC GGG TGG CGC TCG CTC CAG CGG CCG GCG	153
Phe Ala Ala Ala Arg Trp Pro Gly Trp Arg Ser Leu Gln Arg Pro Ala	
20 25 30 35	
CGG CGG AGC GGG CGG GGC GGC GGT GGC GCG GCC CCG GGA CCG TAT CCC	201
Arg Arg Ser Gly Arg Gly Gly Gly Gly Ala Ala Pro Gly Pro Tyr Pro	
40 45 50	
TCC GCC GCC CCT CCC CCG CCC GGC CCC GGC CCC CCT CCC TCC CGG CAG	249
Ser Ala Ala Pro Pro Pro Pro Gly Pro Gly Pro Pro Pro Ser Arg Gln	
55 60 65	
AGC TCG CCT CCC TCC GCC TCA GAC TGT TTT GGT AGC AAC GGC AAC GGC	297
Ser Ser Pro Pro Ser Ala Ser Asp Cys Phe Gly Ser Asn Gly Asn Gly	
70 75 80	
GGC GGC GCG TTT CGG CCC GGC TCC CGG CGG CTC CTT GGT CTC GGC GGC	345
Gly Gly Ala Phe Arg Pro Gly Ser Arg Arg Leu Leu Gly Leu Gly Gly	

85	90	95	
CCT CCC CGC CCC TTC GTC GTC GTC CTT CTC CCC CTC GCC AGC CCG GGC	393		
Pro Pro Arg Pro Phe Val Val Val Leu Leu Pro Leu Ala Ser Pro Gly			
100	105	110	115
GCC CCT CCG GCC GCG CCA ACC CGC GCC TCC CCG CTC GGC GCC CGT GCG	441		
Ala Pro Pro Ala Ala Pro Thr Arg Ala Ser Pro Leu Gly Ala Arg Ala			
120	125	130	
TCC CCG CCG CGT TCC GGC GTC TCC TTG GCG CGC CCG GCT CCC GGC TGT	489		
Ser Pro Pro Arg Ser Gly Val Ser Leu Ala Arg Pro Ala Pro Gly Cys			
135	140	145	
CCC CGC CCG GCG TGC GAG CCG GTG TAT GGG CCC CTC ACC ATG TCG CTG	537		
Pro Arg Pro Ala Cys Glu Pro Val Tyr Gly Pro Leu Thr Met Ser Leu			
150	155	160	
AAG CCC CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CAA	585		
Lys Pro Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln			
165	170	175	
CAG CAG CAG CAG CAG CAG CAG CAG CAG CAG CCG CCG CCC GCG GCT GCC AAT	633		
Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Pro Pro Pro Ala Ala Ala Asn			
180	185	190	195
GTC CGC AAG CCC GGC GGC AGC GGC CTT CTA GCG TCG CCC GCC GCC GCG	681		
Val Arg Lys Pro Gly Gly Ser Gly Leu Leu Ala Ser Pro Ala Ala Ala			
200	205	210	
CCT TCG CCG TCC TCG TCC TCG GTC TCC TCG TCC TCG GCC ACG GCT CCC	729		
Pro Ser Pro Ser Ser Ser Ser Val Ser Ser Ser Ser Ala Thr Ala Pro			
215	220	225	
TCC TCG GTG GTC GCG GCG ACC TCC GGC GGC GGG AGG CCC GGC CTG GGC	777		
Ser Ser Val Val Ala Ala Thr Ser Gly Gly Gly Arg Pro Gly Leu Gly			
230	235	240	

AGA GGT CGA AAC AGT AAC AAA GGA CTG CCT CAG TCT ACG ATT TCT TTT	825
Arg Gly Arg Asn Ser Asn Lys Gly Leu Pro Gln Ser Thr Ile Ser Phe	
245 250 255	
GAT GGA ATC TAT GCA AAT ATG AGG ATG GTT CAT ATA CTT ACA TCA GTT	873
Asp Gly Ile Tyr Ala Asn Met Arg Met Val His Ile Leu Thr Ser Val	
260 265 270 275	
GTT GGC TCC AAA TGT GAA GTA CAA GTG AAA AAT GGA GGT ATA TAT GAA	921
Val Gly Ser Lys Cys Glu Val Gln Val Lys Asn Gly Gly Ile Tyr Glu	
280 285 290	
GGA GTT TTT AAA ACT TAC AGT CCG AAG TGT GAT TTG GTA CTT GAT GCC	969
Gly Val Phe Lys Thr Tyr Ser Pro Lys Cys Asp Leu Val Leu Asp Ala	
295 300 305	
GCA CAT GAG AAA AGT ACA GAA TCC AGT TCG GGG CCG AAA CGT GAA GAA	1017
Ala His Glu Lys Ser Thr Glu Ser Ser Ser Gly Pro Lys Arg Glu Glu	
310 315 320	
ATA ATG GAG AGT ATT TTG TTC AAA TGT TCA GAC TTT GTT GTG GTA CAG	1065
Ile Met Glu Ser Ile Leu Phe Lys Cys Ser Asp Phe Val Val Val Gln	
325 330 335	
TTT AAA GAT ATG GAC TCC AGT TAT GCA AAA AGA GAT GCT TTT ACT GAC	1113
Phe Lys Asp Met Asp Ser Ser Tyr Ala Lys Arg Asp Ala Phe Thr Asp	
340 345 350 355	
TCT GCT ATC AGT GCT AAA GTG AAT GGC GAA CAC AAA GAG AAG GAC CTG	1161
Ser Ala Ile Ser Ala Lys Val Asn Gly Glu His Lys Glu Lys Asp Leu	
360 365 370	
GAG CCC TGG GAT GCA GGT GAA CTC ACA GCC AAT GAG GAA CTT GAG GCT	1209
Glu Pro Trp Asp Ala Gly Glu Leu Thr Ala Asn Glu Glu Leu Glu Ala	
375 380 385	
TTG GAA AAT GAC GTA TCT AAT GGA TGG GAT CCC AAT GAT ATG TTT CGA	1257

Leu Glu Asn Asp Val Ser Asn Gly Trp Asp Pro Asn Asp Met Phe Arg
 390 395 400
 TAT AAT GAA GAA AAT TAT GGT GTA GTG TCT ACG TAT GAT AGC AGT TTA 1305
 Tyr Asn Glu Glu Asn Tyr Gly Val Val Ser Thr Tyr Asp Ser Ser Leu
 405 410 415
 TCT TCG TAT ACA GTG CCC TTA GAA AGA GAT AAC TCA GAA GAA TTT TTA 1353
 Ser Ser Tyr Thr Val Pro Leu Glu Arg Asp Asn Ser Glu Glu Phe Leu
 420 425 430 435
 AAA CGG GAA GCA AGG GCA AAC CAG TTA GCA GAA GAA ATT GAG TCA AGT 1401
 Lys Arg Glu Ala Arg Ala Asn Gln Leu Ala Glu Glu Ile Glu Ser Ser
 440 445 450
 GCC CAG TAC AAA GCT CGA GTG GCC CTG GAA AAC GAT GAT AGG AGT GAG 1449
 Ala Gln Tyr Lys Ala Arg Val Ala Leu Glu Asn Asp Asp Arg Ser Glu
 455 460 465
 GAA GAA AAA TAC ACA GCA GTT CAG AGA AAT TCC AGT GAA CGT GAG GGG 1497
 Glu Glu Lys Tyr Thr Ala Val Gln Arg Asn Ser Ser Glu Arg Glu Gly
 470 475 480
 CAC AGC ATA AAC ACT AGG GAA AAT AAA TAT ATT CCT CCT GGA CAA AGA 1545
 His Ser Ile Asn Thr Arg Glu Asn Lys Tyr Ile Pro Pro Gly Gln Arg
 485 490 495
 AAT AGA GAA GTC ATA TCC TGG GGA AGT GGG AGA CAG AAT TCA CCG CGT 1593
 Asn Arg Glu Val Ile Ser Trp Gly Ser Gly Arg Gln Asn Ser Pro Arg
 500 505 510 515
 ATG GGC CAG CCT GGA TCG GGC TCC ATG CCA TCA AGA TCC ACT TCT CAC 1641
 Met Gly Gln Pro Gly Ser Gly Ser Met Pro Ser Arg Ser Thr Ser His
 520 525 530
 ACT TCA GAT TTC AAC CCG AAT TCT GGT TCA GAC CAA AGA GTA GTT AAT 1689
 Thr Ser Asp Phe Asn Pro Asn Ser Gly Ser Asp Gln Arg Val Val Asn

535	540	545	
GGA GGT GTT CCC TGG CCA TCG CCT TGC CCA TCT CCT TCC TCT CGC CCA	1737		
Gly Gly Val Pro Trp Pro Ser Pro Cys Pro Ser Pro Ser Ser Arg Pro			
550	555	560	
CCT TCT CGC TAC CAG TCA GGT CCC AAC TCT CTT CCA CCT CGG GCA GCC	1785		
Pro Ser Arg Tyr Gln Ser Gly Pro Asn Ser Leu Pro Pro Arg Ala Ala			
565	570	575	
ACC CCT ACA CGG CCG CCC TCC AGG CCC CCC TCG CGG CCA TCC AGA CCC	1833		
Thr Pro Thr Arg Pro Pro Ser Arg Pro Pro Ser Arg Pro Ser Arg Pro			
580	585	590	595
CCG TCT CAC CCC TCT GCT CAT GGT TCT CCA GCT CCT GTC TCT ACT ATG	1881		
Pro Ser His Pro Ser Ala His Gly Ser Pro Ala Pro Val Ser Thr Met			
600	605	610	
CCT AAA CGC ATG TCT TCA GAA GGG CCT CCA AGG ATG TCC CCA AAG GCC	1929		
Pro Lys Arg Met Ser Ser Glu Gly Pro Pro Arg Met Ser Pro Lys Ala			
615	620	625	
CAG CGA CAT CCT CGA AAT CAC AGA GTT TCT GCT GGG AGG GGT TCC ATA	1977		
Gln Arg His Pro Arg Asn His Arg Val Ser Ala Gly Arg Gly Ser Ile			
630	635	640	
TCC AGT GGC CTA GAA TTT GTA TCC CAC AAC CCA CCC AGT GAA GCA GCT	2025		
Ser Ser Gly Leu Glu Phe Val Ser His Asn Pro Pro Ser Glu Ala Ala			
645	650	655	
ACT CCT CCA GTA GCA AGG ACC AGT CCC TCG GGG GGA ACG TGG TCA TCA	2073		
Thr Pro Pro Val Ala Arg Thr Ser Pro Ser Gly Gly Thr Trp Ser Ser			
660	665	670	675
GTG GTC AGT GGG GTT CCA AGA TTA TCC CCT AAA ACT CAT AGA CCC AGG	2121		
Val Val Ser Gly Val Pro Arg Leu Ser Pro Lys Thr His Arg Pro Arg			
680	685	690	

TCT CCC AGA CAG AAC AGT ATT GGA AAT ACC CCC AGT GGG CCA GTT CTT	2169
Ser Pro Arg Gln Asn Ser Ile Gly Asn Thr Pro Ser Gly Pro Val Leu	
695 700 705	
GCT TCT CCC CAA GCT GGT ATT ATT CCA ACT GAA GCT GTT GCC ATG CCT	2217
Ala Ser Pro Gln Ala Gly Ile Ile Pro Thr Glu Ala Val Ala Met Pro	
710 715 720	
ATT CCA GCT GCA TCT CCT ACG CCT GCT AGT CCT GCA TCG AAC AGA GCT	2265
Ile Pro Ala Ala Ser Pro Thr Pro Ala Ser Pro Ala Ser Asn Arg Ala	
725 730 735	
GTT ACC CCT TCT AGT GAG GCT AAA GAT TCC AGG CTT CAA GAT CAG AGG	2313
Val Thr Pro Ser Ser Glu Ala Lys Asp Ser Arg Leu Gln Asp Gln Arg	
740 745 750 755	
CAG AAC TCT CCT GCA GGG AAT AAA GAA AAT ATT AAA CCC AAT GAA ACA	2361
Gln Asn Ser Pro Ala Gly Asn Lys Glu Asn Ile Lys Pro Asn Glu Thr	
760 765 770	
TCA CCT AGC TTC TCA AAA GCT GAA AAC AAA GGT ATA TCA CCA GTT GTT	2409
Ser Pro Ser Phe Ser Lys Ala Glu Asn Lys Gly Ile Ser Pro Val Val	
775 780 785	
TCT GAA CAT AGA AAA CAG ATT GAT GAT TTA AAG AAA TTT AAG AAT GAT	2457
Ser Glu His Arg Lys Gln Ile Asp Asp Leu Lys Lys Phe Lys Asn Asp	
790 795 800	
TTT AGG TTA CAG CCA AGT TCT ACT TCT GAA TCT ATG GAT CAA CTA CTA	2505
Phe Arg Leu Gln Pro Ser Ser Thr Ser Glu Ser Met Asp Gln Leu Leu	
805 810 815	
AAC AAA AAT AGA GAG GGA GAA AAA TCA AGA GAT TTG ATC AAA GAC AAA	2553
Asn Lys Asn Arg Glu Gly Glu Lys Ser Arg Asp Leu Ile Lys Asp Lys	
820 825 830 835	
ATT GAA CCA AGT GCT AAG GAT TCT TTC ATT GAA AAT AGC AGC AGC AAC	2601

Ile	Glu	Pro	Ser	Ala	Lys	Asp	Ser	Phe	Ile	Glu	Asn	Ser	Ser	Ser	Asn	
				840					845					850		
TGT	ACC	AGT	GGC	AGC	AGC	AAG	CCG	AAT	AGC	CCC	AGC	ATT	TCC	CCT	TCA	2649
Cys	Thr	Ser	Gly	Ser	Ser	Lys	Pro	Asn	Ser	Pro	Ser	Ile	Ser	Pro	Ser	
				855				860						865		
ATA	CTT	AGT	AAC	ACG	GAG	CAC	AAG	AGG	GGA	CCT	GAG	GTC	ACT	TCC	CAA	2697
Ile	Leu	Ser	Asn	Thr	Glu	His	Lys	Arg	Gly	Pro	Glu	Val	Thr	Ser	Gln	
				870				875						880		
GGG	GTT	CAG	ACT	TCC	AGC	CCA	GCA	TGT	AAA	CAA	GAG	AAA	GAC	GAT	AAG	2745
Gly	Val	Gln	Thr	Ser	Ser	Pro	Ala	Cys	Lys	Gln	Glu	Lys	Asp	Asp	Lys	
				885				890						895		
GAA	GAG	AAG	AAA	GAC	GCA	GCT	GAG	CAA	GTT	AGG	AAA	TCA	ACA	TTG	AAT	2793
Glu	Glu	Lys	Lys	Asp	Ala	Ala	Glu	Gln	Val	Arg	Lys	Ser	Thr	Leu	Asn	
900					905				910					915		
CCC	AAT	GCA	AAG	GAG	TTC	AAC	CCA	CGT	TCC	TTC	TCT	CAG	CCA	AAG	CCT	2841
Pro	Asn	Ala	Lys	Glu	Phe	Asn	Pro	Arg	Ser	Phe	Ser	Gln	Pro	Lys	Pro	
					920				925					930		
TCT	ACT	ACC	CCA	ACT	TCA	CCT	CGG	CCT	CAA	GCA	CAA	CCT	AGC	CCA	TCT	2889
Ser	Thr	Thr	Pro	Thr	Ser	Pro	Arg	Pro	Gln	Ala	Gln	Pro	Ser	Pro	Ser	
					935				940					945		
ATG	GTG	GGT	CAT	CAA	CAG	CCA	ACT	CCA	GTT	TAT	ACT	CAG	CCT	GTT	TGT	2937
Met	Val	Gly	His	Gln	Gln	Pro	Thr	Pro	Val	Tyr	Thr	Gln	Pro	Val	Cys	
				950				955						960		
TTT	GCA	CCA	AAT	ATG	ATG	TAT	CCA	GTC	CCA	GTG	AGC	CCA	GGC	GTG	CAA	2985
Phe	Ala	Pro	Asn	Met	Met	Tyr	Pro	Val	Pro	Val	Ser	Pro	Gly	Val	Gln	
				965				970						975		
CCT	TTA	TAC	CCA	ATA	CCT	ATG	ACG	CCC	ATG	CCA	GTG	AAT	CAA	GCC	AAG	3033
Pro	Leu	Tyr	Pro	Ile	Pro	Met	Thr	Pro	Met	Pro	Val	Asn	Gln	Ala	Lys	

980	985	990	995	
ACA TAT AGA GCA GTA CCA AAT ATG CCC CAA CAG CGG CAA GAC CAG CAT	3081			
Thr Tyr Arg Ala Val Pro Asn Met Pro Gln Gln Arg Gln Asp Gln His				
1000	1005	1010		
CAT CAG AGT GCC ATG ATG CAC CCA GCG TCA GCA GCG GGC CCA CCG ATT	3129			
His Gln Ser Ala Met Met His Pro Ala Ser Ala Ala Gly Pro Pro Ile				
1015	1020	1025		
GCA GCC ACC CCA CCA GCT TAC TCC ACG CAA TAT GTT GCC TAC AGT CCT	3177			
Ala Ala Thr Pro Pro Ala Tyr Ser Thr Gln Tyr Val Ala Tyr Ser Pro				
1030	1035	1040		
CAG CAG TTC CCA AAT CAG CCC CTT GTT CAG CAT GTG CCA CAT TAT CAG	3225			
Gln Gln Phe Pro Asn Gln Pro Leu Val Gln His Val Pro His Tyr Gln				
1045	1050	1055		
TCT CAG CAT CCT CAT GTC TAT AGT CCT GTA ATA CAG GGT AAT GCT AGA	3273			
Ser Gln His Pro His Val Tyr Ser Pro Val Ile Gln Gly Asn Ala Arg				
1060	1065	1070	1075	
ATG ATG GCA CCA CCA ACA CAC GCC CAG CCT GGT TTA GTA TCT TCT TCA	3321			
Met Met Ala Pro Pro Thr His Ala Gln Pro Gly Leu Val Ser Ser Ser				
1080	1085	1090		
GCA ACT CAG TAC GGG GCT CAT GAG CAG ACG CAT GCG ATG TAT GCA TGT	3369			
Ala Thr Gln Tyr Gly Ala His Glu Gln Thr His Ala Met Tyr Ala Cys				
1095	1100	1105		
CCC AAA TTA CCA TAC AAC AAG GAG ACA AGC CCT TCT TTC TAC TTT GCC	3417			
Pro Lys Leu Pro Tyr Asn Lys Glu Thr Ser Pro Ser Phe Tyr Phe Ala				
1110	1115	1120		
ATT TCC ACG GGC TCC CTT GCT CAG CAG TAT GCG CAC CCT AAC GCT ACC	3465			
Ile Ser Thr Gly Ser Leu Ala Gln Gln Tyr Ala His Pro Asn Ala Thr				
1125	1130	1135		

CTG CAC CCA CAT ACT CCA CAC CCT CAG CCT TCA GCT ACC CCC ACT GGA 3513
 Leu His Pro His Thr Pro His Pro Gln Pro Ser Ala Thr Pro Thr Gly
 1140 1145 1150 1155
 CAG CAG CAA AGC CAA CAT GGT GGA AGT CAT CCT GCA CCC AGT CCT GTT 3561
 Gln Gln Gln Ser Gln His Gly Gly Ser His Pro Ala Pro Ser Pro Val
 1160 1165 1170
 CAG CAC CAT CAG CAC CAG GCC GCC CAG GCT CTC CAT CTG GCC AGT CCA 3609
 Gln His His Gln His Gln Ala Ala Gln Ala Leu His Leu Ala Ser Pro
 1175 1180 1185
 CAG CAG CAG TCA GCC ATT TAC CAC GCG GGG CTT GCG CCA ACT CCA CCC 3657
 Gln Gln Gln Ser Ala Ile Tyr His Ala Gly Leu Ala Pro Thr Pro Pro
 1190 1195 1200
 TCC ATG ACA CCT GCC TCC AAC ACG CAG TCG CCA CAG AAT AGT TTC CCA 3705
 Ser Met Thr Pro Ala Ser Asn Thr Gln Ser Pro Gln Asn Ser Phe Pro
 1205 1210 1215
 GCA GCA CAA CAG ACT GTC TTT ACG ATC CAT CCT TCT CAC GTT CAG CCG 3753
 Ala Ala Gln Gln Thr Val Phe Thr Ile His Pro Ser His Val Gln Pro
 1220 1225 1230 1235
 GCG TAT ACC AAC CCA CCC CAC ATG GCC CAC GTA CCT CAG GCT CAT GTA 3801
 Ala Tyr Thr Asn Pro Pro His Met Ala His Val Pro Gln Ala His Val
 1240 1245 1250
 CAG TCA GGA ATG GTT CCT TCT CAT CCA ACT GCC CAT GCG CCA ATG ATG 3849
 Gln Ser Gly Met Val Pro Ser His Pro Thr Ala His Ala Pro Met Met
 1255 1260 1265
 CTA ATG ACG ACA CAG CCA CCC GGC GGT CCC CAG GCC GCC CTC GCT CAA 3897
 Leu Met Thr Thr Gln Pro Pro Gly Gly Pro Gln Ala Ala Leu Ala Gln
 1270 1275 1280
 AGT GCA CTA CAG CCC ATT CCA GTC TCG ACA ACA GCG CAT TTC CCC TAT 3945

Ser Ala Leu Gln Pro Ile Pro Val Ser Thr Thr Ala His Phe Pro Tyr

1285

1290

1295

ATG ACG CAC CCT TCA GTA CAA GCC CAC CAC CAA CAG CAG TTG 3987

Met Thr His Pro Ser Val Gln Ala His His Gln Gln Gln Leu

1300

1305

1310

TAAGGCTGCC CTGGAGGAAC CGAAAGGCCA AATTCCCTCC TCCCTTCTAC TGCTTGTACC 4047

AACTGGAAGC ACAGAAACT AGAATTTTCAT TTATTTTGT TTTAAATAT ATATGTTGAT 4107

TTCTTGTAAC ATCCAATAGG AATGCTAACA GTTCACTTGC AGTGGAGAT ACTTGACCG 4167

AGTAGAGGCA TTTAGGAAC TGGGGGCTAT TCCATAATTC CATATGCTGT TTCAGAGTCC 4227

CGCAGGTACC CCAGCTCTGC TTGCCGAAAC TGGAAGTTAT TTATTTTTA ATAACCCTTG 4287

AAAGTCATGA ACACATCAGC TAGCAAAAGA AGTAACAAGA GTGATTCTTG CTGCTATTAC 4347

TGCTAAAAAA AAAAAAAAAA 4367

SEQ ID NO: 2

SEQUENCE LENGTH: 203

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

CACCACCAGC AACAGCAACA GCAGCAGCAG CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG 60

CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG 120

CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG 180

CAGCATCAGC GAAACTCTGG GCC 203

SEQ ID NO: 3

SEQUENCE LENGTH: 20

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

CACCACCAGC AACAGCAACA

20

SEQ ID NO: 4

SEQUENCE LENGTH: 20

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

GGCCCAGAGT TTCCGTGATG

20

SEQ ID NO: 5

SEQUENCE LENGTH: 165

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG 60

CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG 120

CAGCAGCAGC AGCAGCAGCA GCAGCAGCAG CAGCAGCAGC AGCAG 165

SEQ ID NO: 6

SEQUENCE LENGTH: 21

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

CCCTCACCAT GTCGCTGAAG C

21

SEQ ID NO: 7

SEQUENCE LENGTH: 19

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

CGACGCTAGA AGGCCGCTG

19

SEQ ID NO: 8

SEQUENCE LENGTH: 19

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

CTTGCGGACA TTGGCAGCC

19

SEQ ID NO: 9

SEQUENCE LENGTH: 27

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

TTCTCTCAGC CAAAGCCTTC TACTACC

27

SEQ ID NO: 10

SEQUENCE LENGTH: 19

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

TATCCGCAGC TCCGCTCCC

19

SEQ ID NO: 11

SEQUENCE LENGTH: 20

SEQUENCE TYPE: nucleic acid

STRNDEDNESS: single

TOPOLOGY: linear

SEQUENCE DESCRIPTION

AGCCGGGGCCG AAACGGGCGG

20

CLAIMS

1. A nucleic acid fragment comprising a nucleic acid region encoding an amino acid sequence shown in SEQ ID NO: 1 (provided that the number of repeat units of Gln
5 from the 166th to 188th amino acid varies between 15 and 100).
2. The nucleic acid fragment according to claim 1, wherein said nucleic acid region is the region from 49nt to 3987nt (provided that the number of repeat units of
10 CAG or CAA in the region from the 543nt to 612nt varies between 15 and 100, and that the CAA in this region may be CAG).
3. A protein having the amino acid sequence encoded by said nucleic acid fragment according to claim 1 or 2.
- 15 4. An antibody which undergoes antigen-antibody reaction with said protein according to claim 3.
5. An antisense nucleic acid having a size of not less than 15 bp, which hybridizes with the mRNA transcribed from the nucleic acid fragment according to claim 1 or 2
20 so as to inhibit translation thereof.
6. A recombinant vector comprising said nucleic acid fragment according to claim 1 or 2 incorporated into an expression vector which can express a desired gene in human body, which recombinant vector can express said
25 nucleic acid fragment in human body.
7. A method comprising introducing said recombinant vector according to claim 6 into human body and

expressing said nucleic acid fragment according to claim
1 or 2 in said human body.

ABSTRACT

A sequence-determined cDNA fragment of the causative gene of SCA2 is disclosed. The cDNA fragment according to the present invention comprises a nucleic acid region
5 encoding an amino acid sequence shown in SEQ ID NO: 1
(provided that the number of repeat units of Gln from the 166th to 188th amino acid varies between 15 and 100).

1/6

1 TATCCGCACCTCCGCTCCACCCGGCGCCTCGGCGCGCCCGCCCTCCGATGCGCTCAGCG
 1 F-1006 M R S A
 61 GCCGCAGCTCCTCGGAGTCCCGCGGTGGCCACCGAGTCTCGCCGCTTCGCCGCAGCCAGG
 5 A A A P R S P A V A T E S R R F A A A R
 121 TGGCCCCGGGTGGCGCTCGCTCCAGCGGCCGGCGCGGCGGAGCGGGCGGGGCGGCGGTGGC
 25 W P G W R S L Q R P A R R S G R G G G G
 181 GCGGCCCCGGGACCGTATCCCTCCGCGCCCTCCCCCGCCCGGCCCCGGCCCCCTCCC
 45 A A P G P Y P S A A P P P P G P G P P P
 241 TCCCGGCAGAGCTCGCCTCCCTCCGCCTCAGACTGTTTTGGTAGCAACGGCAACGGCGGC
 65 S R Q S S P P S A S D C F G S N G N G G
 301 GCGCGTTTCGGCCCCGGCTCCCGCGGCTCCTTGGTCTCGGCGGGCCTCCCCGCCCTTC
 85 R-1002 G A F R P G S R R L L G L G G P P R P F
 361 GTCGTCGTCCTTCTCCCCCTCGCCAGCCCGGGCGCCCTCCGCGCGGCCAACCCGCGCC
 105 V V V L L P L A S P G A P P A A P T R A
 421 TCCCCGCTCGGCGCCCGTGGCTCCCCCGCGGTTCCGCGGTCTCCTTGGCGCGCCCGGT
 125 S P L G A R A S P P R S G V S L A R P A
 481 CCCGGCTGTCCCCGCCCGCGGTGCGAGCCGGTGTATGGGCCCTCACCATGTCGCTGAAG
 145 P G C P R P A C E P V Y G F-1 P L T M S L K
 541 CCCCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAGCAACAGCAGCAGCAGCAG
 165 P Q
 601 CAGCAGCAGCAGCCGCCGCCCGCGGCTGCCAATGTCCGCAAGCCCCGGCGGCAGCGCCTT
 185 Q Q Q Q P P P A A A N V R K P G G S G L
 661 CTAGCGTCGCCCGCGCGCGCCTTCGCCGTCCTCGTCTCGGTCTCCTCGTCTCGGCC
 205 R-1 L A S P A A A P S P S S S S V S S S S A
 721 ACGGCTCCCTCCTCGGTGGTGGCGGCGACCTCCGGCGGCGGAGGCCCGGCTGGGCAGA
 225 T A P S S V V A A T S G G G R P G L G R
 781 GGTCAAACAGTAACAAAGGACTGCCTCAGTCTACGATTTCTTTTGATGGAATCTATGCA
 245 G R N S N K G L P Q S T I S F D G I Y A
 841 AATATGAGGATGGTTCATATACTTACATCAGTTGTTGGCTCCAAATGTGAAGTACAAGTG
 265 N M R M V H I L T S V V G S K C E V Q V
 901 AAAAATGGAGGTATATATGAAGGAGTTTTTAAACTTACAGTCCGAAGTGTGATTTGGTA
 285 K N G G I Y E G V F K T Y S P K C D L V
 961 CTTGATGCCGCACATGAGAAAAGTACAGAATCCAGTTCGGGGCCGAAACGTGAAGAAATA
 305 L D A A H E K S T E S S S G P K R E E I
 1021 ATGGAGAGTATTTTGTTCAAATGTTTACAGTCTTGTGTGGTACAGTTTAAAGATATGGAC
 325 M E S I L F K C S D F V V V Q F K D M D
 1081 TCCAGTTATGCAAAAAGAGATGCTTTTACTGACTCTGCTATCAGTGCTAAAGTGAATGGC
 345 S S Y A K R D A F T D S A I S A K V N G
 1141 GAACACAAAGAGAAGGACCTGGAGCCCTGGGATGCAGGTGAACACAGCCAATGAGGAA
 365 E H K E K D L E P W D A G E L T A N E E
 1201 CTTGAGGCTTTGGAAAATGACGTATCTAATGGATGGGATCCCAATGATATGTTTCGATAT
 385 L E A L E N D V S N G W D P N D M F R Y
 1261 AATGAAGAAAATTATGGTGTAGTGTCTACGTATGATAGCAGTTTATCTTCGTATACAGTG
 405 N E E N Y G V V S T Y D S S L S S Y T V

Fig. 1

1321 CCCTTAGAAAGAGATAA ACTCAGAAGAATTTT TAAAACGGGAAGCAAGGGCAAACCAGTTA
425 P L E R D N S E E F L K R E A R A N Q L
1381 GCAGAAGAAATTTGAGTCAAGTGCC CAGTACAAAGCTCGAGTGGCCCTGGAAAACGATGAT
445 A E E I E S S A Q Y K A R V A L E N D D
1441 AGGAGTGAGGAAGAAAAATACACAGCAGTT CAGAGAAATTCAGTGAACGTGAGGGGCAC
465 R S E E E K Y T A V Q R N S S E R E G H
1501 AGCATAAACACTAGGGAAAATAAATATATTCCTCTCGGACAAAGAAATAGAGAAGTCATA
485 S I N T R E N K Y I P P G Q R N R E V I
1561 TCCTGGGGAAGTGGGAGACAGAATTCACCGCGTATGGGCCAGCCTGGATCGGGCTCCATG
505 S W G S G R Q N S P R M G Q P G S G S M
1621 CCATCAAGATCCACTTCTCACACTTCAGATTTCAACCCGAATTCTGGTTCAGACCAAAGA
525 P S R S T S H T S D F N P N S G S D Q R
1681 GTAGTTAATGGAGGTGTTCCCTGGCCATCGCCTTGCCCATCTCCTTCTCGCCCCACCT
545 V V N G G V P W P S P C P S P S S R P P
1741 TCTCGCTACCAGTCAGGTCCCAACTCTCTTCCACCTCGGGCAGCCACCCCTACACGGCCG
565 S R Y Q S G P N S L P P R A A T P T R P
1801 CCCTCCAGGCCCCCTCGCGGCCATCCAGACCCCGTCTCACCCCTCTGCTCATGGTTCT
585 P S R P P S R P S R P P S H P S A H G S
1861 CCAGCTCTGTCTCTACTATGCCTAAACGCATGTCTTCAGAAGGGCCTCCAAGGATGTCC
605 P A P V S T M P K R M S S E G P P R M S
1921 CCAAAGGCCCAGCGACATCCTCGAAATCACAGAGTTTCTGCTGGGAGGGGTTCATATCC
625 P K A Q R H P R N H R V S A G R G S I S
1981 AGTGGCCTAGAATTTGTATCCCAACCCACCCAGTGAAGCAGCTACTCCTCCAGTAGCA
645 S G L E F V S H N P P S E A A T P P V A
2041 AGGACCAGTCCCTCGGGGGGAACGTGGTCATCAGTGGTTCAGTGGGGTTCCAAGATTATCC
665 R T S P S G G T W S S V V S G V P R L S
2101 CCTAAAACTCATAGACCCAGGTCTCCAGACAGAACAGTATTGGAAATACCCCCAGTGGG
685 P K T H R P R S P R Q N S I G N T P S G
2161 CCAGTTCTTGCTTCTCCCCAAGCTGGTATTATTCCAAGTGAAGCTGTTGCCATGCCTATT
705 P V L A S P Q A G I I P T E A V A M P I
2221 CCAGCTGCATCTCCTACGCCTGCTAGTCCTGCATCGAACAGAGCTGTTACCCCTTCTAGT
725 P A A S P T P A S P A S N R A V T P S S
2281 GAGGCTAAAGATTCCAGGCTTCAAGATCAGAGGCAGAACTCTCCTGCAGGGAATAAAGAA
745 E A K D S R L Q D Q R Q N S P A G N K E
2341 AATATTAACCCAATGAAACATCACCTAGCTTCTCAAAAGCTGAAAACAAAGGTATATCA
765 N I K P N E T S P S F S K A E N K G I S
2401 CCAGTTGTTTCTGAACATAGAAAACAGATTGATGATTTAAAGAAATTTAAGAATGATTTT
785 P V V S E H R K Q I D D L K K F K N D F
2461 AGGTTACAGCCAAGTTCTACTTCTGAATCTATGGATCAACTACTAAACAAAAATAGAGAG
805 R L Q P S S T S E S M D Q L L N K N R E
2521 GGAGAAAAATCAAGAGATTTGATCAAAGACAAAATGAACCAAGTGCTAAGGATTCTTTT
825 G E K S R D L I K D K I E P S A K D S F
2581 ATTGAAAAATAGCAGCAGCAACTGTACCAGTGGCAGCAGCAAGCCGAATAGCCCCAGCATT
845 I E N S S S N C T S G S S K P N S P S I

Fig. 2

2641 TCCCCTTCAATACTTAGTAACACGGAGCACAAGAGGGGACCTGAGGTCACTTCCCAAGGG
 865 S P S I L S N T E H K R G P E V T S Q G
 2701 GTTCAGACTTCCAGCCCAGCATGTAAACAAGAGAAAGACGATAAGGAAGAGAAGAAAGAC
 885 V Q T S S P A C K Q E K D D K E E K K D
 2761 GCAGCTGAGCAAGTTAGGAAATCAACATTGAATCCCAATGCAAAGGAGTTCAACCCACGT
 905 A A E Q V R K S T L N P N A K E F N P R
 2821 TCCTTCTCTCAGCCAAAGCCTTCTACTACCCCACTTCACCTCGGCCTCAAGCACAACTT
 F-13
 925 S F S Q P K P S T T P T S P R P Q A Q P
 2881 AGCCCATCTATGGTGGGTCAACAGCCAACCTCCAGTTTATACTCAGCCTGTTTGT
 945 S P S M V G H Q Q P T P V Y T Q P V C F
 2941 GCACCAAATATGATGTATCCAGTCCCAGTGAGCCCAGGCGTGCAACCTTTATACCCAATA
 965 A P N M M Y P V P V S P G V Q P L Y P I
 3001 CCTATGACGCCCATGCCAGTGAATCAAGCCAAGACATATAGAGCAGTACCAAATATGCC
 985 P M T P M P V N Q A K T Y R A V P N M P
 3061 CAACAGCGGCAAGACCAGCATCATCAGAGTGCCATGATGCACCCAGCGTCAGCAGCGGGC
 1005 Q Q R Q D Q H H Q S A M M H P A S A A G
 3121 CCACCGATTGCAGCCACCCACCAGCTTACTCCACGCAATATGTTGCCTACAGTCCTCAG
 1025 P P I A A T P P A Y S T Q Y V A Y S P Q
 3181 CAGTTCCTCAAATCAGCCCTTGTTCAGCATGTGCCACATTATCAGTCTCAGCATCCTCAT
 1045 Q F P N Q P L V Q H V P H Y Q S Q H P H
 3241 GTCTATAGTCCTGTAATACAGGGTAATGCTAGAATGATGGCACCACCAACACACGCCCAG
 1065 V Y S P V I Q G N A R M M A P P T H A Q
 3301 CCTGGTTTATGATCTTCTTTCAGCAACTCAGTACGGGGCTCATGAGCAGACGCATGCGATG
 1085 P G L V S S S A T Q Y G A H E Q T H A M
 3361 TATGCATGTCCCAAATTACCATAACAAGAGACAAGCCCTTCTTTCTACTTTGCCATT
 1105 Y A C P K L P Y N K E T S P S F Y F A I
 3421 TCCACGGGCTCCCTTGCTCAGCAGTATGCGCACCTAACGCTACCCTGCACCCACATACT
 1125 S T G S L A Q Q Y A H P N A T L H P H T
 3481 CCACACCCTCAGCCTTCAGCTACCCCACTGGACAGCAGCAAAGCCAACATGGTGGAGT
 1145 P H P Q P S A T P T G Q Q Q S Q H G G S
 3541 CATCTGCACCCAGTCCTGTTTCAGCACCATCAGCACCAGGCCGCCAGGCTCTCCATCTG
 1165 H P A P S P V Q H H Q H Q A A Q A L H L
 3601 GCCAGTCCACAGCAGCAGTCAGCCATTTACCACGCGGGGCTTGCGCCAACTCCACCCTCC
 1185 A S P Q Q Q S A I Y H A G L A P T P P S
 3661 ATGACACCTGCCTCCAACAGCAGTCGCCACAGAATAGTTTCCCAGCAGCACAACTCAGACT
 1205 M T P A S N T Q S P Q N S F P A A Q Q T
 3721 GTCTTTACGATCCATCCTTCTCAGTTTCAGCCGCGGTATACCAACCCACCCACATGGCC
 1225 V F T I H P S H V Q P A Y T N P P H M A
 3781 CACGTACCTCAGGCTCATGTACAGTCAGGAATGGTTCCTTCTCATCCAACTGCCCATGCG
 1245 H V P Q A H V Q S G M V P S H P T A H A
 3841 CCAATGATGCTAATGACGACACAGCCACCCGGCGGTCCCCAGGCCGCCCTCGCTCAAAGT
 1265 P M M L M T T Q P P G G P Q A A L A Q S
 3901 GCACTACAGCCCATTCAGTCTCGACAACAGCGCATTTCCCTATATGACGCACCTTCA
 1285 A L Q P I P V S T T A H F P Y M T H P S
 3961 GTACAAGCCCACCACCAACAGCAGTTGTAAGGCTGCCCTGGAGGAACCGAAAGGCCAAAT
 1305 V Q A H H Q Q Q L *

Fig. 3

4/6

4021 TCCCTCCTCCCTTCTACTGCTTCTACCAACTGGAAGCACAGAAAAGTAGAATTTTCATTTA
 4081 TTTTGTTTTTTAAAATATATATGTTGATTTCTTGTAACATCCAATAGGAATGCTAACAGTT
 4141 CACTTGCAGTGGAAGATACTTGGACCGAGTAGAGGCATTTAGGAACTTGGGGGCTATTCC
 4201 ATAATTCCATATGCTGTTTTAGAGTCCCGCAGGTACCCCGAGCTCTGCTTGCCGAACTGG
 4261 AAGTTATTTATTTTTTTAATAACCCCTGAAAGTCATGAACACATCAGCTAGCAAAAGAAGT
 4321 AACAAAGAGTGATTCTTGCTGCTATTACTGCT(A)_n

Fig. 4




Fig. 5




Fig. 6

6/6




Fig. 7

COMBINED DECLARATION AND POWER OF ATTORNEY
FOR PATENT APPLICATION

As a below named inventor, I hereby declare that: my residence post office address and citizenship are as stated next to my name; that I verily believe that I am the original, first and sole inventor (if only one inventor is named below) or a joint inventor (if plural inventors are named below) of the subject matter which is claimed and for which a patent is sought on the invention entitled: ***"cDNA Fragment of Causative Gene of Spinocerebellar Ataxia Type 2"**

the specification of which is attached hereto unless one of the following boxes is checked:

☐ The Specification was filed on

and was assigned Serial No.

and was amended on

☒ was filed as PCT international application number **PCT/JP97/03946** on **October 30, 1997** and was amended under PCT Article 19 on (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56.

I do not know and do not believe the same was ever known or used in the United States of America before my or our invention thereof, or patented or described in any printed publication in any country before my or our invention thereof, or more than one year prior to this application, that the same was not in public use or on sale in the United States of America more than one year prior to this application, that the invention has not been patented or made the subject of an inventor's certificate issued before the date of this application in any country foreign to the United States of America on an application filed by me or my legal representatives or assigns more than twelve months prior to this application, and that no application for patent or inventor's certificate on this invention has been filed in any country foreign to the United States of America prior to this application by me or my legal representatives or assigns, except as follows:

I hereby claim foreign priority benefits under Title 35, United States Code, § 119 of any foreign application(s) for patent or inventor's certificate listed below:

Prior Foreign Application(s)

Priority Claimed

304059/96

Japan

10.30.96

Yes

(Number)

(Country)

(Month/Day/Year Filed)

All Foreign Applications, if any, for any Patent or Inventor's Certificate Filed More Than 12 Months Prior To The Filing Date of This Application:

Country

Application No.

Date of Filing (Month/Day/Year)

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, § 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)

(Filing Date)

(Status-patented,
pending, abandoned)

I hereby appoint the following attorneys to prosecute this application and/or an international application based on this application and to transact all business in the Patent and Trademark Office connected therewith and in connection with the resulting patent based on instructions received from the entity who first sent the application papers to the attorneys identified below, unless the inventor(s) or assignee provides said attorneys with a written notice to the contrary:

RAYMOND C. STEWART (Reg. No. 21,066) TERRELL C. BIRCH (Reg. No. 19,382)

JOSEPH A. KOLASCH (Reg. No. 22,463) ANTHONY L. BIRCH (Reg. No. 26,122)

JAMES M. SLATTERY (Reg. No. 28,380) BERNARD L. SWEENEY (Reg. No. 24,448)

DONALD C. KOLASCH (Reg. No. 23,038) MICHAEL K. MUTTER (Reg. No. 29,680)

CHARLES GORENSTEIN (Reg. No. 29,271) GERALD M. MURPHY, JR.

LEONARD R. SVENSSON (Reg. No. 30,330) (Reg. No. 28,977)

MARC S. WEINER (Reg. No. 32,181) TERRY L. CLARK (Reg. No. 32,644)

ANDREW D. MEIKLE (Reg. No. 32,868)

Send Correspondence to: BIRCH, STEWART, KOLASCH AND BIRCH

8110 Gatehouse Road

Suite 500 East

P.O. Box 747

Falls Church, Virginia 22040-0747

Telephone: (703)205-8000

Facsimile: (703)205-8050

RECEIVED SEP 14 1996

14

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	*Date
<u>Shoji</u>	<u>TSUJI</u>	<i>Shoji Tsuji</i>	June 23, 1998
RESIDENCE(City, State & Country)			CITIZENSHIP
<u>Niigata-shi, Niigata, JAPAN</u>			Japanese
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
<u>1-10-45, Kobari, Niigata-shi, Niigata 950-21 JAPAN</u>			
GIVEN NAME	FAMILY NAME	INVENTOR'S SIGNATURE	*Date
<u>Kazuhiro</u>	<u>SANPEI</u>	<i>Kazuhiro Sanpei</i>	June 23, 1998
RESIDENCE(City, State & Country)			CITIZENSHIP
<u>Niigata-shi, Niigata, JAPAN</u>			Japanese
POST OFFICE ADDRESS (Complete Street Address including City, State & Country)			
<u>A-203 Advantage, 98-1, Oshima, Niigata-shi, Niigata 950 JAPAN</u>			

JPX

JPX